

13. Berechne ohne Taschenrechner!

$$(a) \quad {}^3\log \frac{1}{\sqrt[5]{9}} = x \quad 3^x = 9^{-\frac{1}{5}} \\ 3^x = (3^2)^{-\frac{1}{5}} = 3^{-\frac{2}{5}} \\ x = -\frac{2}{5}$$

$$(b) \quad {}^5\log \sqrt[4]{125} = x \quad 5^x = (5^3)^{\frac{1}{4}} \\ 5^x = 5^{\frac{3}{4}} \\ x = \frac{3}{4}$$

$$(c) \quad \lg \sqrt{0,1} = x \quad 10^x = 10^{-\frac{1}{2}} \\ {}^{10}\log \left(\frac{1}{10} \right)^{\frac{1}{2}} = x \quad x = -\frac{1}{2}$$

$$(d) \quad \lg \frac{1}{\sqrt[3]{100}} = x \quad 10^x = 10^{-\frac{2}{3}} \\ {}^{10}\log 10^{-\frac{2}{3}} = x \quad x = -\frac{2}{3}$$

$$(e) \quad \ln \sqrt[5]{\frac{1}{e^2}} = x \quad e^x = e^{-\frac{2}{5}} \\ {}^e\log e^{-\frac{2}{5}} = x \quad x = -\frac{2}{5}$$

$$(f) \quad \ln \frac{e}{\sqrt{e}} = x \quad e^x = e^{\frac{1}{2}} \\ {}^e\log e^{1-\frac{1}{2}} = x \quad x = \frac{1}{2} \\ {}^e\log e^{\frac{1}{2}} = x$$

$$(g) \quad {}^a\log 1 = x \quad a^x = 1 \\ x = 0$$

$$(h) \quad {}^2\log -4 = x \quad 2^x = -4 \\ \text{nicht möglich!}$$

$$(i) \quad \ln 1 = x \quad e^x = 1 \\ {}^e\log 1 = x \quad x = 0$$

$$(j) \quad \ln e = x \quad e^x = e \\ {}^e\log e = x \quad x = 1$$

14. Berechne x!

(a) ${}_x \log 9 = -2$

$$x^{-2} = 9$$

$$\frac{1}{x^2} = 9 \quad | \cdot x^2 | : 9$$

$$x^2 = \frac{1}{9}$$

$$x = \frac{1}{3} \text{ (nur positiv, da log nur}$$

für $\mathbf{R}^+$ definiert ist)

(b) ${}_x \log a^4 = 2$

$$x^2 = a^4 \quad | \sqrt{}$$

$$x = a^2$$

(c) ${}_x \log 27 = -\frac{3}{4}$

$$x^{-\frac{3}{4}} = 27$$

$$\frac{1}{\sqrt[4]{x^3}} = 27 \quad |^4 | \sqrt[3]{}$$

$$\frac{1}{x} = 3^4 \quad | \cdot x | : 3^4$$

$$x = \frac{1}{81}$$

(d) ${}_x \log 5 = 2$

$$x^2 = 5 \quad | \sqrt{}$$

$$x = \sqrt{5}$$

(e) ${}_4 \log x = -1$

$$4^{-1} = x$$

$$x = \frac{1}{4}$$

(f) $\lg x = -\frac{1}{4}$

$${}_{10} \log x = -\frac{1}{4}$$

$$10^{-\frac{1}{4}} = x$$

$$x = \frac{1}{\sqrt[4]{10}}$$

(g) ${}_a \log x = a$

$$x = a^a$$

15. Berechne!

(a) $\lg \left(\underbrace{\lg 1}_1 0 \right) = \lg 1 = 0$

$${}_{10} \log 10 = x$$

$$10^x = 10$$

$$x = 1$$

$${}_{10} \log 1 = y$$

$$10^y = 1$$

$$y = 0$$

(b) $\lg \left(\underbrace{\ln e}_{10}^{10} \right) = \lg 10 = 1$

$${}_e \log e^{10} = x$$

$$e^x = e^{10}$$

$$x = 10$$

$${}_{10} \log 10 = y$$

$$10^y = 10$$

$$y = 1$$

(c) $\ln \left(\underbrace{\ln e}_1 \right) = \ln 1 = 0$

$${}_e \log e = x$$

$$e^x = e$$

$$x = 1$$

$${}_e \log 1 = y$$

$$e^y = 1$$

$$y = 0$$

(d) $\ln \left(\underbrace{\lg 1}_e 0^e \right) = \ln e = 1$

$${}_{10} \log 10^e = x$$

$$10^x = 10^e$$

$$x = e$$

$${}_e \log e = y$$

$$e^y = e$$

$$y = 1$$

16. Zerlege mit Hilfe der Logarithmusregeln!

$$\begin{aligned} \text{(a)} \quad \log \left(x^2 \sqrt{\frac{y}{\sqrt[3]{z}}} \right) &= \log x^2 + \log \left(\frac{y}{\sqrt[3]{z}} \right)^{\frac{1}{2}} = \\ &= 2 \log x + \frac{1}{2} \log \left(\frac{y}{\sqrt[3]{z}} \right) = \\ &= 2 \log x + \frac{1}{2} \left(\log y - \frac{1}{3} \log z \right) = \\ &= 2 \log x + \frac{1}{2} \log y - \frac{1}{6} \log z \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad \log \frac{\sqrt[4]{2a} \cdot \sqrt[3]{b^2}}{4b \cdot \sqrt{a}} &= \log \sqrt[4]{2a} + \log \sqrt[3]{b^2} - \log 4 - \log b - \log \sqrt{a} = \\ &= \frac{1}{4} \log 2a + \frac{1}{3} \log b^2 - \log 2^2 - \log b - \frac{1}{2} \log a = \\ &= \frac{1}{4} \log 2 + \frac{1}{4} \log a + \frac{2}{3} \log b - 2 \log 2 - \log b - \frac{1}{2} \log a = \\ &= -\frac{7}{4} \log 2 - \frac{1}{4} \log a - \frac{1}{3} \log b \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad \log \left(4x^5 \sqrt[3]{\frac{x^2 - a^2}{2x^3 \sqrt{a}}} \right) &= \log 4 + \log x^5 + \log \left(\frac{x^2 - a^2}{2x^3 \sqrt{a}} \right)^{\frac{1}{3}} = \\ &= \log 2^2 + \log x^5 + \frac{1}{3} \log \left(\frac{x^2 - a^2}{2x^3 \sqrt{a}} \right) = \\ &= 2 \log 2 + 5 \log x + \frac{1}{3} \left(\log(x^2 - a^2) - \log(2x^3 \sqrt{a}) \right) = \\ &= 2 \log 2 + 5 \log x + \frac{1}{3} \left(\log((x+a)(x-a)) - \log 2 - \log x^3 - \log \sqrt{a} \right) = \\ &= 2 \log 2 + 5 \log x + \frac{1}{3} \left(\log(x+a) + \log(x-a) - \log 2 - 3 \log x - \frac{1}{2} \log a \right) = \\ &= 2 \log 2 + 5 \log x + \frac{1}{3} \log(x+a) + \frac{1}{3} \log(x-a) - \frac{1}{3} \log 2 - \log x - \frac{1}{6} \log a = \\ &= \frac{5}{3} \log 2 + 4 \log x + \frac{1}{3} \log(x+a) + \frac{1}{3} \log(x-a) - \frac{1}{6} \log a \end{aligned}$$

17. Stelle als Logarithmus eines einzigen Terms dar!

$$\begin{aligned}
 \text{(a)} \quad & \frac{1}{3}[5 \log x + 2 \log(x-y)] - \log x - \frac{1}{2} \log y = \\
 & = \frac{1}{3}[\log x^5 + \log(x-y)^2] - \log x - \log \sqrt{y} = \\
 & = \frac{1}{3}[\log(x^5 \cdot (x-y)^2)] - \log(x \cdot \sqrt{y}) = \\
 & = \log \frac{\sqrt[3]{x^5 \cdot (x-y)^2}}{x \cdot \sqrt{y}}
 \end{aligned}$$

$$\begin{aligned}
 \text{(b)} \quad & \frac{1}{2} \left[8 \log x + \frac{3}{4} \left(5 \log x - \frac{1}{3} \log c + 4 \log a + \log b - \frac{2}{3} \log 2 \right) \right] = \\
 & = \frac{1}{2} \left[\log x^8 + \frac{3}{4} (\log x^5 - \log \sqrt[3]{c} + \log a^4 + \log b - \log \sqrt[3]{2^2}) \right] = \\
 & = \frac{1}{2} \left[\log x^8 + \frac{3}{4} \log \frac{a^4 b x^5}{\sqrt[3]{2^2 \sqrt[3]{c}}} \right] = \frac{1}{2} \left[\log x^8 + \log \sqrt[4]{\left(\frac{a^4 b x^5}{\sqrt[3]{2^2 \sqrt[3]{c}}} \right)^3} \right] = \\
 & = \frac{1}{2} \left[\log x^8 + \log \sqrt[4]{\left(\frac{a^4 b x^5}{\sqrt[3]{2^2 \sqrt[3]{c}}} \right)^3} \right] = \\
 & = \log \sqrt{x^8 \sqrt[4]{\left(\frac{a^4 b x^5}{\sqrt[3]{2^2 \sqrt[3]{c}}} \right)^3}}
 \end{aligned}$$

$$\begin{aligned}
 \text{(c)} \quad & 3 \log x - \frac{2}{3} [4 \log y + 3 \log z] + \frac{1}{2} \left[\frac{1}{3} \log x - \log y \right] = \\
 & = \log x^3 - \frac{2}{3} [\log y^4 + \log z^3] + \frac{1}{2} [\log \sqrt[3]{x} - \log y] = \\
 & = \log x^3 - \frac{2}{3} \log y^4 z^3 + \frac{1}{2} \log \frac{\sqrt[3]{x}}{y} = \\
 & = \log x^3 - \log \sqrt[3]{(y^4 z^3)^2} + \log \sqrt{\frac{\sqrt[3]{x}}{y}} = \\
 & = \log \frac{x^3 \sqrt{\frac{\sqrt[3]{x}}{y}}}{\sqrt[3]{(y^4 z^3)^2}}
 \end{aligned}$$

$$\begin{aligned}
 \text{(d)} \quad & 2 \log x + \frac{1}{3} \log y + \log(1-x^2) - 4 \log x + \frac{1}{2} \log x - [\log(1-x) + \log(1+x) - 2 \log x] \\
 & = \\
 & = \log x^2 + \log \sqrt[3]{y} + \log(1-x^2) - \log x^4 + \log \sqrt{x} - \log(1-x) - \log(1+x) + \log x^2 = \\
 & = \log \frac{x^2 \cdot \sqrt[3]{y} \cdot (1-x^2) \cdot \sqrt{x} \cdot x^2}{x^4 \cdot (1-x) \cdot (1+x)} = \\
 & = \log \frac{\sqrt[3]{y} \cdot (1-x^2) \cdot \sqrt{x}}{(1-x^2)} = \\
 & = \log \sqrt{x} \sqrt[3]{y}
 \end{aligned}$$